

## Recent distribution of the Euro-Siberian-sub-Mediterranean species *Elatine alsinastrum* L. (Elatinaceae)

AGNIESZKA POPIELA<sup>1\*</sup>, ANDRZEJ ŁYSKO<sup>2</sup>, ATTILA MOLNÁR<sup>3</sup>

<sup>1</sup> Department of Botany and Nature Conservation, University of Szczecin, Felczaka 3c, 71-412 Szczecin, Poland

<sup>2</sup> Department of Environmental Protection and Management, Western Pomeranian University of Technology, Szczecin, Poland

<sup>3</sup> Department of Botany, University of Debrecen, Egyetem sq. 1, 4032 Hungary

**Abstract** – The general distribution of the endangered Euro-Siberian sub-Mediterranean species *Elatine alsinastrum* L. is provided using literature, web-sources and herbaria dataset. The distribution pattern shows some regularities: occurrence of locations along river valleys, formation of concentrated site clusters in some lowlands, wide distances between locations or site clusters or single locations between their clusters. The distribution patterns in central Europe seem to be rather well related to the history of the human migration in Europe at least since the Late Holocene. The scattered locations on the eastern part of the distribution area are likely to be a consequence of missing information, rather than to the fragmentation of its distribution.

**Keywords:** *Elatine alsinastrum*, ephemeral, distribution

### Introduction

The genus *Elatine* L. (*Malpighiales* Mart., *Elatinaceae* Dumort.) includes approximately 15–25 species occurring in temperate areas of both hemispheres, mostly in the northern. Besides *Elatine*, the family includes the genus *Bergia* L. (TUCKER 1986). Species of the genus *Elatine* are phytogeographically quite interesting, since they are found in all continents, except Antarctica, mainly in the temperate zones, and reach their greatest spread in the northern hemisphere, both in terms of species-diversity and number of locations. Despite this wide distribution, the species of *Elatine* occur rarely, and their sites are usually quite distant from each other (often more than 100 km). They are semi-aquatic therophytes adapted to live in periodically flooded environments. The numbers of populations and

\* Corresponding author, e-mail: popiela@univ.szczecin.pl

Copyright® 2013 by Acta Botanica Croatica, the Faculty of Science, University of Zagreb. All rights reserved.

plants within a population usually show remarkable yearly fluctuations. The plants, usually characterized by an outstanding phenotypic variability and plasticity, have a very short reproduction period and may produce, when environmental conditions are favourable, many seeds, thus preserving the long lasting germination capacity (DEIL 2005; POPIELA and ŁYSKO 2010, 2011; POPIELA et al. 2010, 2011), as KASAHARA et al. (1967) that has been demonstrated for *Elatine triandra* Schkuhr (more than 50 years).

*Elatine alsinastrum* L. is taxonomically isolated within the family Elatinaceae, being the only taxon included in the subgenus *Potamopithys* (Adanson) Seub., and clearly differing from other species of the genus by the presence of leaves arranged in whorls and most frequently a longer and sturdier stem. Other species, with opposite leaves, are comprised in the subgenus *Elatine* (= *Hydropiper* Moesz.) with the sections: *Triandra* Seub. (= *Crypta* (Nutt.) Seub.] and *Elatinella* Seub. (TUCKER 1986). Accordingly, *E. alsinastrum* is relatively easy to identify and both herbarium materials and literature data can be considered reliable. It includes upright amphibian herbs, with stem erect or ascending, mostly simple; leaves in whorls of (4)8–17 (submerged) and of 3 (aerial); flowers usually 3 at a node, mostly sessile or short-stalked; sepals 4, petals 4 and stamens 8. Seeds cylindrical, 0.6–0.9 mm long, 0.2–0.3 mm width, almost straight to slightly bent (0–55°), pits conspicuous, (15–)17–21(–24) in each row.

The species occurs in two forms: the first is terrestrial, and it can be found in wet depressions, while the second is aquatic and can be found in shallow stagnant waters (30–50 cm deep), in flooded rice fields and meadows, pond waters and on their verges, lake shores, in periodically dry puddles and water bodies, drainage ditches, in wet depressions in cultivated fields and on wet roadsides. It frequently occurs with other ephemeral wetland species, such as *E. triandra* Schkurb., *Limosella aquatica* L., *Lindernia procumbens* (Krock.) Philcox, *Cyperus fuscus* L., *Plantago major* L. s.l., *Gnaphalium uliginosum* L., *Eleocharis acicularis* (L.) Roem. et Shult., *Peplis portula* Schrnk. (class *Isoëto-Nanojuncetea* Br.-Bl. et Tüxen ex Br.-Bl. et all. 1952, associations: *Myosotido siculae-Isoëtetum velatae* Pottier-Alapetite 1952, *Eleocharito-Limoselletum aquatica* Philippi 1968, *Lindernio-Eleocharitetum ovatae* Pietsch 1973, *Eleocharito acicularis-Schoenoplectetum supine* (Horvatić 1931) Sao et Ubrizsy in Ubrizsy 1951 (= *Elatini-Lindernietum procumbentis* Ubrizsy 1961), *Elatini alsinastri-Juncetum tenageiae* Libbert 1932, *Ranunculo lateriflori-Limoselletum aquatica* Pop 1968 (LIBBERT 1932; OBERDORFER 1957; PIETSCH 1973a, b; POTTIER-ALAPETITE 1952; BRULLO et MINISSALE 1998).

*Elatine alsinastrum* is an Euro-Siberian species, quite rare in the whole area of its distribution range (MEUSEL et al. 1978, HULTÉN and FRIES 1986, UOTILA 2009a). The aim of this work is to provide the current distribution of this endangered taxon.

## Material and methods

This paper is based on both analysis of floristic literature, distribution atlases, web-sources and examination of the specimens kept in 31 herbaria, 20 of them accessed by web. (Tab. 1). A total of 993 floristic data (700 from the literature and herbaria, 293 from the Web) are analyzed, priority being given to literature data over Websources. The latter were used only if considered to be reliable (e.g. if they did not deviate from the range or had an author and exact location or were delivered by herbarium servers). In consequence, 392 web data were removed.

Data were manually mapped using a satellite base image from Google Earth. Calibration, rectification to UTM grid and digitalization were done. The vector layers were prepared by conversion to WGS84 cartographic grid. Electronic data containing geographic co-ordinates were exported to SHP format. When geographic coordinates were missing, records were converted into text format. If detailed descriptions are missing and only cartographic grid locations are given, spatial queries connecting grid area and map were used. All analyses and maps were made using PostGIS database extension software and QGIS (Quantum GIS) application software working in the LINUX environment.

## Results


*Elatine alsinastrum* is a Euro-Siberian sub-Mediterranean species (Fig. 1). The analysis of the current distribution of *E. alsinastrum* allows us to infer that its distribution centre is in the lowland and upland areas of Europe situated to the north and west of Alps and Carpathians. The pattern of its distribution has the form of smaller and larger site clusters which are frequently far away from each other, even many kilometers. They are mainly located along the valleys of the major rivers. The highest concentration of sites was found in

**Tab. 1.** Literature, herbaria and web sources. Herbarium acronyms follow Index Herbariorum (THIERS 2012), but the herbarium full name is given when not listed there.


Area	Data
Asia, North and Eastern Europe (Russia, Kazakhstan, Armenia, Georgia, Ukraine, Moldova, Finland, Lithuania, Belarus)	ADYLOV 1983, CZEREPAKOV 1995, GAGNIDZE 2005, GEIDEMAN 1986, GROSSHEIM 1962, KORSHINSKY 1898, KRYLOV 1939, KUUSK et al. 1996, MAEVSKY 1954, PAPCZENKOV 2006, PAVLOV 1963, PESCHKOVA 2006, PROKUDIN 1987, 1999, SHISHKIN and BOBROV 1974, SHISHKIN 1961, SMALLHAUSEN 1895, TAKHTAJYAN 1966, TICHOMIROVA 1986, TZVELEV 1996, UOTILA 2009b, VEDENKOV 1998, VOLOBAIEV 1991. Specimens found in the following herbaria: K*, B
Central, Southern and South-Eastern Europe (Poland, Czech Republic, Slovakia, Austria, Hungary, Romania, the Balkans, Italy)	BARINA and PIFKÓ 2007, BERGMEIER & ABRAHAMCZYK 2008, BLEČIĆ 1972, BOROS and VAJDA 1957, CESATI et al. 1868–86, DANIN 2004, DAVIS 1967, DESFAYES 2008, DIMITROVA 2009, DOSTÁL and ČERVENKA 1991, DOSTÁL 1989, FARKAS 1999, FINTHA 1994, FIORI and PALEOTTI 1896, GORI et al. 1998, GRDINIĆ et al. 2001, HAYEK 1927, HEJNÝ S. and SLAVÍK 1990, JAKAB 2005, JUHÁSZ-NAGY 1959, MOESZ 1908, MOLNÁR and GULYÁS 2001, MOLNÁR and PFEIFFER 1999, NIKOLIĆ and TOPIĆ 2005, NIKOLIĆ 1994, PEEV 1984, PETKOVIĆ et al. 1998, PIGNATTI 1982, PINKE et al. 2006, POPIELA 2001, POPIELA et al. 2012a, POPIELA et al. 2010, PROHÁZKA and KØISA 1999, SARICA-HATZINIKOLAOU et al. 1994, 1997, SÂVULESCU 1955, SOÓ and MÁTHÉ 1938, STEVANOVIĆ 1999, TRINAJSTIĆ 1975, VLADIMIROV et al. 2010, Specimens found in the following herbaria: K*, B, BP, DE, SAMU, Kazinczy Ferenc Museum, Sátoraljaújhely (Hungary); University of Szeged, Szeged (Hungary); Szent István University, Gödöllő (Hungary); Kazinczy Ferenc Museum, Sátoraljaújhely (Hungary). Biodiversity occurrence data were provided by the University Museums of Norway.

**Tab. 1.** – continued

Area	Data
Western and South-Western Europe and Northern Africa (France, Germany, Switzerland, Spain, Portugal, Morocco, Algeria, Tunisia)	<p>ANONYMUS 2005, ANTONETTI et al. 2006, BENKERT et al. 1996, BOLOMIER CIRUIANO and VELAYOS 1993, COUTINHO 1939, DIARD 2005, FENNANE et al. 1998, FORTUNE 2003, FRANCO 1971, FUKAREK and HENKER 2006, GAMISANS and JEANMONOD, 1993, GRENIER 1992, GUINOCHE et VILMORIN 1982, HAEUPLER and SCHÖNFELDER 1989, HAMMADA et al. 2004, JAHANDIEZ and MAIRE 1932, JEANMONOD and GAMISANS 2007, LE FLOC'H and BOULEZ 2008, LUNAIS et al. 1986, MULLER 2006, NETIEN 1993, POTTIER-ALAPETITE 1979, QUEZEL and SANTA 1962–1963, SAINT-LAGER 1873, SCHOTSMAN and OSSERDET 1967, TOURLET 1908, WOHLGEMUTH 1993.</p> <p>Specimens were found in the following herbaria: K*, B.</p> <p>Biodiversity occurrence data were provided by the following Herbaria: ABH; BC; CIBIO; COFC; E; FCO; GLM; HSS; IBF; Leiner-Herbar Konstanz; LUX; MA; MAF; MGC; OHN; SALA; SANT; SEV; STR; STU.</p> <p>Occurrence data were provided by Information System of the plants of Spain, Real Jardín Botánico, CSIC – Fundación Biodiversidad; retrieved November 20, 2011 from <a href="http://www.anthos.es">www.anthos.es</a>.</p> <p>Biodiversity occurrence data were provided by following database: Aranzadi Zientzi Elkartea; Botany (UPS); Bundesamt für Naturschutz/ Netzwerk Phytodiversitaet Deutschland; Cartografía de vegetación a escala de detalle 1:10.000 de la masa forestal de Andalucía; Flora exsiccata Bavarica; Florabank1; Inventaire national du Patrimoine naturel (INPN); Observations du Conservatoire botanique national du Bassin parisien.; Phanerogamie; Phytochorologie des départements français; Sistema de Información de la vegetación Ibérica y Macaronésica</p>

**Fig. 1.** Distribution area of *Elatine alsinastrum* L.

the French plains (Armorican Massif, Paris, and Aquitaine Basin), in the central zones of the Loire and Seine river valleys, in the upper part of the Rhine river valley, in the East German lowlands and in the western area of Poland, in the Oder and Vistula river valleys (Fig. 2).



**Fig. 2.** Distribution area of *Elatine alsinastrum* L. in central Europe.

Smaller site clusters are in western and central Europe, in the Bohemian Massif, in the Danube and Tisza river valleys, and in the Pannonian Basin. Furthermore, it is found in scattered locations in southern Europe and at the southern limit of its distribution range in single locations in Morocco, Algeria, Tunisia and Palestine. The distribution in eastern Europe (the area of the Vistula-eastern Carpathian Mountains) appears to be unclear (Fig. 1), probably because there is a scarcity of data on detailed locations from this area. Most of the Russian floras have scarce or no information about the occurrence of the taxon at issue or the available information is rather scattered. The analysis of floras of the Baltic countries and information from Belarus show that this species either occurs very rarely there or is absent. In Ukraine, it is found scattered over the whole territory, except the Crimea. There are only scattered locations of *E. alsinastrum* in the European part of Russia from the North and the South by the Volga and the Kama rivers. Site clusters of this taxon are also observed in the Stavropol Krai and Georgia and Armenia where it reaches the southern limit of its range in eastern Europe. In Siberia, single locations are found in the valleys of left tributaries of the Tobol River (the Iset and Tura rivers), in the Irtysh river valley, and in Kazakhstan. The largest number of observations in Siberia has come from natural locations: swamps, flooded meadows, lake shores, flooded lowlands, river meanders and the shores of drying up water bodies.

## Discussion

On the basis of fossil data and molecular findings, DAVIS et al. (2001) have raised a hypothesis that families of the order *Malpighiales* originate in the northern areas of South America, with subsequent migration to North America and later, via the Atlantic, to the Old World. This would suggest that the genus *Elatine* in the northern hemisphere has a history going back at least to the Early Tertiary and floras of that period (e.g. the Mediterranean Tertiary and the Madrean Tertiary), while the present range of the genus originated in the

Holocene. *E. alsinastrum* is the species with the most isolated taxonomic position within the family Elatinaceae; accordingly, a question arises about the formation of its range in the Holocene, the refuge for this species during the Pleistocene glaciations and the migration routes.

The distribution area of *E. alsinastrum* L. shows a remarkable geographic disparity and certain regularities: the formation of concentrated site clusters in some lowlands, and large distances between locations or site clusters or single locations between their clusters. The situation in central Europe appears to be well known, with smaller and larger site clusters (often along river valleys) which are frequently far away from each other, even many kilometers away. So, disjunctions inside the range seem to occur, indicating that the spreading of diaspores does not occur easily despite good habitat conditions. It is possible that the species is a relic of the former natural wetland flora, its sites thus able to remain permanent for a very long time (soil seed bank) and the distribution pattern is a reflection of history of colonization in Europe at least since the Late Holocene. Many locations have been changed by human beings and thus, it seems that the range of *E. alsinastrum* may be also extended anthropogenically, with human-created habitats; however, the species is also spread by birds. Dispersal of diaspores by water and marsh birds (epi- and/or endozoochoria) is possible, although documentation in the genus *Elatine* is only known for one species (KERNER 1895).

In a discussion of methodological problems in mapping the range of *E. alsinastrum* differences in the quality of data and the methods being used for their preparation should be emphasized. In this study, the universal WGS 84 coordinate system was used, while the maps employed are made in local or regional coordinate systems. The best method is calibration of maps in the system, in which they were made, followed by their vectorisation and reduction to a uniform coordinate system. Using this method, data do not lose quality when the map is being scaled up, while a point on it always preserves the specific scale (see also: POPIELA et al. 2012b).

The comparison of the maps resulting from the present study with those published earlier (MEUSEL et al. 1978, HULTÉN and FRIES 1986) shows a larger amount of data (also with reference to those available in the last twenty five years) and a greater precision, which was allowed by the mapping method adopted. Changes in the distribution of *E. alsinastrum* relate mainly to central Spain, and southern France. The method adopted has also allowed verification of the dispersal character of the distribution of the species in central Europe.

### Acknowledgements

The authors are very grateful to Gábor Sramkó for linguistic improvements. The work was provided to A. Popiela by a Ministry, Department of Scientific Research in Warsaw (Poland) grant Nr PO4C03525. We wish to thank the keepers of the herbaria for access to the material and to anonymous reviewers for giving valuable comments.

### References

- ADYLOV, T. A., 1983: *Conspectus Floraes Asiae Mediae*, 7 (In Russian). USSR Academy of Sciences/Uzbek Academy of Sciences, Moscow-Tashkent.

- ANONYMUS, 2005: Catalogue raisonné des plantes vasculaires de la Gironde. Mémoires de la Société Linnéenne de Bordeaux 4, Bordeaux.
- ANTONETTI, P., BRUGEL, E., KESSLE, F., 2006: Atlas de la flore d'Auvergne. Conservatoire botanique du Massif central, Chavaniac-Lafayette.
- BARINA, Z., PIFKÓ, D., 2007: Botanical research in the Visegrád Mountains [In Hungarian]. Kitaibelia 12, 9–25.
- BENKERT, D., FUKAREK, F., KORSCH, H. (eds.), 1996: Verbreitungsatlas der Farn- und Blütenpflanzen Ostdeutschlands. Gustav Fischer, Jena.
- BERGMEIER, E., ABRAHAMCZYK, S., 2008: Current and historical diversity and new records of wetland plants in Crete, Greece. *Wilddenowia* 38, 433–453.
- BLEČIĆ, V., 1972: Elatinaceae. In: JOSIFOVIĆ, M. (ed.), Flore de la République Socialiste de Serbie, 3, 126–128. Académie Serbe des sciences et des arts, Belgrad.
- BOLOMIER, A. C., CATTIN, P., 1999: La flore du département de l'Ain, inventaire complet. Connaissance de la flore de l'Ain, Bourg en Bresse.
- BOREAU, A., 1849: Flore du centre de la France et du bassin de la Loire. Paris.
- BOROS, Á., VAJDA, L., 1957: The Sphagnum-bogs of Bakony and Balaton-region [In Hungarian]. *Annales Instituti Biologici Hungaricae Academiae Scientiarum (Tihany)* 24, 282–287.
- BOUDIN, L., CORDIER, J., MORET, J., 2007: Atlas de la flore remarquable du Val de Loire. Muséum National d'Histoire Naturelle, Paris.
- BRULLO, S., MINISSALE, P. 1998. Considerazioni sintassonomiche sulla classe Isoeto-Nano-juncetea. *Itineraria Geobotanica* 11, 263–290.
- BUGNON, F., FELZINES, J. C., GOUX, R., LOISEAU, J. E., ROYER, J. M., 1998: Nouvelle Flore de Bourgogne, 3. Atlas de répartition, clés des groupements végétaux et suppléments aux tomes I1, 2. *Bulletin Scientifique de Bourgogne*, Dijon.
- CESATI, V., PASSERINI, G., GIBELLI, G., 1868–1886: Compendio della flora Italiana. F. Val-ladi, Milano.
- CIRUJANO, S., VELAYOS, M., 1993: *Elatine* L. In: CASTROVIEJO, S. (ed.), Flora Iberica, 3. Real Jardín Botánico, Consejo Superior de Investigaciones Científicas, Madrid.
- COUTINHO, A. X. P., 1939: Flora de Portugal: plantas vasculares, 2. Bertrand, Lisboa.
- CZEREPOV, S. K., 1995: Vascular plants of Russia and adjacent states (the former USSR). Cambridge University Press, Cambridge.
- DANIN, A., 2004: Distribution atlas of plants in the flora Palestina area. The Israel Academy of Sciences and Humanities, Jerusalem.
- DAVIS P. H. F., 1967: Flora of Turkey and the Aegean Island. Edinburgh University Press; Edinburgh.
- DAVIS, C. C., ANDERSON, W. R., DONOGUE, M. J., 2001. Phylogeny of Malpighiaceae: evidence from chloroplast *ndhF* and *trnL-F* nucleotide sequences. *American Journal of Botany* 88, 1830–1846.
- DEIL, U., 2005: A review on habitats, plant traits and vegetation of ephemeral wetlands – a global perspective. *Phytocoenologia* 35, 533–705.

- DESFAYES, M., 2008: Flore vasculaire herbacée des eaux douces et des milieux humides de la Sardaigne. *Flora Mediterranea* 18, 247–331.
- DIARD, L., 2005: *Atlas de La Flore d'Ille-et-Vilaine*. Siloë edition, Nantes.
- DIMITROVA, D., 2009: *Elatine alsinastrum*. In: PETROVA, A., VLADIMIROV, V. (eds.), Red list of Bulgarian vascular plants. *Phytologica Balcanica* 15, 68.
- DOSTÁL, J., 1989: New flora of Czechoslovakia 1, 2 (In Czech]. Academia, Praha.
- DOSTÁL, J., ČERVENKA, M., 1991: Key for higher plants (In Slovakian], I. Slovenske Pedagogicke Nakladatel'stvo, Bratislava.
- FARKAS, S., 1999: Protected plants of Hungary (In Hungarian). Mezogazda Kiadó, Budapest.
- FENNANE, M., IBN TATTOU, M. 1998: Catalogue des plantes vasculaires rares, menacées ou endémiques du Maroc. *Bocconeia* 8, 1–243.
- FENNANE, M., IBN TATTOU, M., MATHEZ, J., OUYAHYA, A., EL OULAIDI, J., 1999: Flore pratique du Maroc, 1. Travaux de l'Institut Scientifique Rabat Série Botanique 36, 288.
- FINTHA, I., 1994: The vascular plants of Észak-Alföld region (In Hungarian). A KTM természetvédelmi Hivatalának tanulmánykötetei I. TermészetBúvár Alapítvány Kiadó, Budapest.
- FIORI, A., PALEOTTI, G., 1896: Flora analitica d'Italia. Tipografia del Seminario, Padova.
- FORTUNE, C., 2003: *Damasonium alisma* Mill. et *Elatine alsinastrum* L. dans le marais de Gannedel (Ille-et-Vilaine). *Erica* 17, 3–14.
- FRANCO, J. A., 1971: Nova Flora de Portugal. I. Edição do Autor, Lisboa.
- FUKAREK, F., HENKER, H., 2006: Flora von Mecklenburg-Vorpommern. Weissdornverlag, Jena.
- GAGNIDZE, R., 2005: Vascular plants of Georgia. A nomenclatural checklist. Georgia Academy of Sciences. N. Ketshoveli Institute of Botany, Tbilisi.
- GAMISANS, J., JEANMONOD, D., 1993: Catalogue des plantes vasculaires de la Corse. In: JEANMONOD, D., BURDET, H. M. (eds.), Complements au Prodrome de la Flore de la Corse. Annexe nr 3. Conservatoire et Jardin botaniques Ville de Genève, Genève.
- GEIDEMAN, T. S., 1986: Identification key of higher plants of Moldavian Soviet Socialist Republic (In Russian]. Shtiintsa, Kishinev.
- GORI, S., LAKATOS, G., ARADI, C., KISS, M., BITSKEY, K., 1998: The vegetation of Meggyes-marsh in the initial stage of rehabilitation (In Hungarian). *Kitaibelia* 3,103–104.
- GRDINIĆ, B., BOŽA, P. P., KNEŽEVIĆ, A., KRIVOKUĆIN, I., 2001: Overview of the flora of Western Backa region (In Croatian). Norma 7, 137–157.
- GRENIER, E., 1992: Flore d'Auvergne. Société Linnéenne de Lyon, Lyon.
- GROSSHEIM, A. A., 1962: Flora of the Caucasus 6 (In Russian). Acad. Sci. USSR, Moscow-Leningrad.
- GUINOCHET, M., VILMORIN DE, R., 1982: Flore de France, 4. Centre National de la Recherche Scientifique, Paris.
- HAEUPLER, H., SCHÖNFELDER, P., 1989: *Atlas der Farn- und Blütenpflanzen der BR Deutschland*, 2. Ulmer, Stuttgart.

- HAMMADA, S., DAKKI, M., IBN TATTOU, M., OUYAHYA, A., FENNANE, M., 2004: Analyse de la biodiversité floristique des zones humides du Maroc. Flore rare, menacée et halophile. *Acta Botanica Malacitana* 29, 43–66.
- HAYEK, A., 1927: *Prodromus Florae Peninsulae Balcanicae*, 1. Dahlem, Berlin.
- HEJNÝ, S., SLAVÍK, B., 1990: Flora of the Czech Republic 2 (In Czech). Academia, Praha.
- HULTÉN, E., FRIES, M., 1986: Atlas of North European vascular plants, north of the tropic of Cancer, 1. Koeltz Scientific Books, Königstein.
- JAHANDIEZ, E., MAIRE, R., 1932: Catalogue des plantes du Maroc, 2. Imprimerie Minerva, Alger.
- JAKAB, G., 2005: New data to the flora of South Tiszántúl region. (In Hungarian). *Flora Pannonica* 3, 91–119.
- JEANMONOD, D., GAMISANS, J., 2007: Flora Corsica. Edisud, Aix-en-Provence.
- JONSELL, B., JARVIS, C. E., 2002: Lectotypifications of Linnaean names for Flora Nordica (Brassicaceae – Apiaceae). *Nordic Journal of Botany* 22, 67–86.
- JUHÁSZ-NAGY, P., 1959: Les associations des prairies et pâtu rages de la plaine »Beregi-sík«. *Acta Universitatis Debreceniensis de Ludovico Kossuth Nominate* 4, 195–228.
- KASAHARA, Y., NISHI, K., UEGAMA, Y., 1967: Studies on the germination of seeds and their growth in rush (*Juncus bufonius* L. var. *decipiens* Buchen.) and weeds, buried for about 50 years. *Hikobia* 5, 91–103.
- KERNER VON MARIALAUN, A., 1895: The natural history of plants. Their forms, growth, reproduction, and distribution, 2. Blackie et Son, London.
- KORSHINSKY, S., 1898: Tentamen florae Rossiae orientalis. Memoires de l' Académie impériale des sciences de St. Petersburg, St. Petersburg, 8 Serie. Classe physic-mathématique 7.
- KRYLOV, P., 1939: Flora of Western Siberia 7 (In Russian). Nauka, Tomsk.
- KUUSK, V., TABAKA, L., YANKYAVICHENE, R., 1996: Flora of the Baltic countries: compendium of vascular plants. Eesti Loodusfoto AS, Tartu.
- LE FLOC'H, E., BOULOS, L., 2008: Flore de Tunisie. Catalogue synonymique commenté. Chez l'auteur, Montpellier.
- LIBBERT, W., 1932: Die Vegetationseinheiten der neumärkischen Plönetal. *Verhandlungen des Botanischen Verein von Berlin und Brandenburg* 78, 72–137.
- LUNAIS, B., MAUBER, P., GUILLOT, G., 1986: Répartition des plantes rares et localisées de Sologne. *Atlas préliminaire 1978–1985: Coll. Inventaires de faune et de flore*, fasc. 31. Secrétariat de la faune et de la flore, Muséum national d'histoire naturelle, Paris.
- MAEVSKY, P. F., 1954: Flora of middle belt of the European part of the USSR (In Russian). Selkhozgiz, Moscow et Leningrad.
- MEUSEL, H., JÄGER, E., RAUSCHERT, S., WEINERT, E., 1978: Vergleichende Chorologie der Zentraleuropäischen Flora, 2. Karten. G. Fischer Verlag, Jena.
- MOESZ, G., 1908: Die Elatinen Ungarns. *Magyar Botanikai Lapok* 7, 2–35.
- MOLNÁR, V. A., GULYÁS, G., 2000: Mapping of the Nanocyperion species on the Great Hungarian Plain in 2000 (In Hungarian). *Kitaibelia* 6, 169–198.

- MOLNÁR, V. A., PFEIFFER, N., 1999: Researching of muddy vegetation in Hungary in year of high floods and inland waters (In Hungarian). *Kitaibelia* 4, 391–421.
- MULLER, S., 2006: Les Plantes protégées de Lorraine: distribution, écologie, conservation, biotope. Parthénope, Mèze.
- NETIEN, G., 1993: Flore lyonnaise. Société Linnéenne de Lyon, Lyon.
- NIKOLIĆ, T., 1994: *Index Florae Croaticae*, 1. *Natura Croatica* 3 (Suppl. 2), 1–116.
- NIKOLIĆ, T., TOPIĆ, J. (eds.), 2005: Red data book of vascular flora of Croatia (In Croatian). Ministerstvo kulture, Državni zavod za zaštitu prirode, Zagreb.
- OBERDORFER, E., 1957: Süddeutsche Pflanzengesellschaften. *Pflanzensoziologie* 10, 1–564.
- PAPCZENKOV, W. G., 2006: *Elatine alsinastrum* L. (In Russian). In: Red data book of Republic of Tatarstan. Idel-Press, Kazan.
- PAVLOV, N. V., 1963: Flora of Kazakhstan, 6 (In Russian). Academia of Science Kazakh SSR, Alma-Ata.
- PEEV, D., 1984: *Elatine alsinastrum* L. In: VELCHEV, V. (ed.), Red data book of the PR Bulgaria, 1. Plants, 241 (In Bulgarian). Publishing House of the Bulgarian Academy of Sciences, Sofia.
- PESCHKOVA, G. A. (ed.), 2006: Flora of Siberia, 10. Geraniaceae–Cornaceae. Science Publishers Inc, Enfield, New Hampshire.
- PETKOVIĆ, B., MARTIN, P., TATIĆ, B., 1998: Genus *Elatine* L. (In Serbian). In the Flora of Serbia. Proceedings 2 Biological Congress of Macedonia, 24.
- PIETSCH, W., 1973a: Beitrag zur Gliederung der europäischen Zwerbinsengesellschaften (Isoëto-Nanojuncetea Br.-Bl. et Tx. 1943). *Vegetatio* 28, 401–438.
- PIETSCH, W., 1973b: Zur Soziologie und Oecologie der Zwerbinsen-Gesellschaften Ungarns (Klasse Isoëto-Nanojucetea Br.-Bl. Et Tx. 1943). *Acta Botanicae Academiae Scientiarum Hungariae* 19, 269–288.
- PIGNATTI, S., 1982: Flora d’Italia. 3. Edagricole, Bologna.
- PINKE, G., PÁL, R., KIRÁLY, G., SZENDRODI, V., 2006: Data to the weed flora in Somogy territory (South-Western Hungary) (In Hungarian). *Botanikai Közlemények* 93, 53–68.
- POPIELA, A., 2001: *Elatine alsinastrum* L. In: ZAJAC, A., ZAJAC, M. (eds.), Distribution atlas of vascular plants in Poland. Nakladem Pracowni Chorologii Komputerowej Instytutu Botaniki Uniwersytetu Jagiellonskiego, Kraków.
- POPIELA, A., ŁYSKO, A., 2010: The distribution of *Elatine macropoda* Guss. (Elatinaceae). *Acta Societatis Botanicorum Poloniae* 79, 81–86.
- POPIELA, A., ŁYSKO, A., 2011: The distribution of species of the section *Elatinella* Seub. (*Elatine* L., Elatinaceae) in Europe. In: ZEMANEK, B. (ed.), Geobotanist and taxonomist. A volume dedicated to professor Adam Zajac on the 70<sup>th</sup> anniversary of his birth, 147–159. Instytut of Botany, Jagiellonian University, Cracow.
- POPIELA, A., ŁYSKO, A., KONOPSKA, K., 2012a: *Elatine alsinastrum* L. (In Polish). In: KAZMIERCZAKOWA, K., ZARZYCKI, Z., MIREK (eds.), Polish red plant book. Ferns and flowering plants (In Polish). Instytut Botaniki im. W. Szafera PAN, Kraków (in press).
- POPIELA, A., ŁYSKO, A., MOLNÁR, V., A., 2012b: The dsitribution of *Elatine hydropiper* L.. *Acta Societatis Botanicorum Poloniae* 81, 137–143.

- POPIELA, A., ŁYSKO, A., NALEPKA, D., 2010: Analysis of distribution patterns of selected ephemeral wetland species in Western Pomerania (NW Poland, NE Germany). *Biodiversity: Research and Conservation* 19, 55–64.
- POPIELA, A., ŁYSKO, A., WIECZOREK, A., NALEPKA, D., 2011: The distribution of *Elatine hexandra* Lapierre (Elatinaceae). *Acta Societatis Botanicorum Poloniae* 80, 27–32.
- POTTIER-ALAPETITE, G., 1952: Note préliminaire sur l'Isoëtion Tunisie. *Bulletin de la Société Botanique de France* 99, 4–6
- POTTIER-ALAPETITE, G., 1979: Flore de la Tunisie. Angiospermes – Dicotyledones. 1. Apetales Dialypetales. Imprimerie Officielle de la République Tunisienne, Tunis.
- PROHÁZKA, F., KRISA, B., 1999: *Elatine alsinastrum* L. (In Czech). In: ČEROVSKÝ, J., FERÁKOVÁ, V., HOLUB, J., MAGLOCKÝ, Š., PROCHÁZKA, F. (eds.), Červená kniha ohrožených druhů rostlin a živočichů ČR a SR. 5, 145. Vyšší rostiny. Priroda a. s., Bratislava.
- PROKUDIN, J. N. (ed.), 1999: Key of higher plants of Ukraine (In Ukrainian]. Naukova Dumka, Ukrainian National Academy of Sciences, Kiev.
- PROKUDIN, J. N. (ed.), 1987: Manual of the higher plants of Ukraine (In Russian). Ukrainian National Academy of Sciences, Kiev.
- QUEZEL, P., SANTA, S., 1962–1963: Nouvelle flore de l'Algérie et des régions désertiques méridionales, 2. Centre National Recherche Scientifique, Paris.
- SAINT-LAGER, J.-B., 1873: Catalogue des plantes vasculaires de la flore du bassin du Rhône. Librairie H. Georg, Lyon.
- SARIKA-HATZINIKOLAOU, M., KOUMLI-SOVANTSI, L., YANNITSAROS, A., 1997: Macrophytes in four Alpine aquatic ecosystems of N. Pindos (Ipiros, Greece). *Phyton* 37, 19–30.
- SARIKA-HATZINIKOLAOU, M., KOUMLI-SOVANTSI, L., YANNITSAROS, A., 1994: *Myriophyllum alterniflorum* DC. (Haloragaceae), a new record for the Greek. *Flora* 34, 243–246
- SĂVULESCU, T. (ed.), 1955: Flora of People Republic of Romania (In Romanian). Academiei Republicii Populare Române, Bucuresti.
- SCHOTSMAN, H. D., BOSSERDET, P., 1966: Notes sur *Elatine brochonii* Clav. *Bulletin du Centre D'Etudes et de Recherches Scientifiques, Biarritz* 6, 251–267.
- SHISHKIN, B. K., BOBROV, E. G., 1974: Flora of the USSR, 15. Israel Program for Scientific Translations, Jerusalem.
- SHISHKIN, B. K., 1961: Flora of Leningrad region, 3 (In Russian). Izdatelstvo Leningradskovo Universiteta, Leningrad.
- SMALLHAUSEN, I., 1895: Flora of middle and south Russia, Krym and Caucasus, 1. N. Oglolin Magazin, Kiev – St. Petersburg.
- SOÓ, R., MÁTHÉ, I., 1938: The Flora of the Tiszántúl region (In Hungarian]. Magyar Flóramuvek 2, Debrecen.
- STEVANOVIĆ, V., 1999: The red data book of flora of Serbia, 1. Extinct and critically endangered taxa. Institute for the Protection of Nature of Serbia, Belgrade.
- TAKHTAJAN, A. L., 1966: Systematics and phylogeny of Magnoliophytorum (In Russian). Nauka, Moscow – Leningrad.

- THIERS, B., 2012: *Index Herbariorum*: a global directory of public herbaria and associated staff, new york botanical gardenžs virtual herbarium. Retrieved October 23, 2011 from <http://sweetgum.nybg.org/ih>
- TIKHOMIROVA, W. I., 1986: Key for plants of Meščery, 1 (In Russian). Izdatelstvo Moskovskogo Universiteta, Moscow.
- TOURLET, E-H., 1908: Catalogue raisonné des Plantes vasculaires du departament d'Indre-et-Loire. Tours, Paris.
- TRINAJSTIĆ, I., 1975: Analytical flora of Yugoslavia 2 (In Croatian). Institut za Botaniku Sveučilišta u Zagrebu, Zagreb.
- TUCKER, G. C., 1986: The genera of Elatinaceae in the southeastern United States. Journal of the Arnold Arboretum 67, 471–483.
- TZVELEV, N. N. (ed.), 1996: Flora of eastern Europe, 9 (In Russian). Mir et Semja, St. Petersburg.
- UOTILA, P. 2009a: Elatinaceae. In: Euro+Med Planbase – the information resource for Euro-Mediterranean plant diversity. Retrieved May 20, 2012 from [www.bgbm.org/EuroPlusMed/PTaxonDetail.asp?/Nemeld=19030etPTReFk=7300000](http://www.bgbm.org/EuroPlusMed/PTaxonDetail.asp?/Nemeld=19030etPTReFk=7300000)
- UOTILA, P., 2009b: *Elatine*. In: Flora Nordica. 6. Retrieved November 23, 2011 from [www.floranordica.org](http://www.floranordica.org)
- VEDENKOV, Y. P., 1998: Specificity of vegetation of the Great Chaply Vale. In: GAVRYLENK, W. C. (ed.), Falz-Fein Biosphere Reserve, 20–25. Ascania Nova, Kherson.
- VLADIMIROV, V., DANE, F., STEVANOVIĆ, V., TAN, K., 2010: New floristic records in the Balkans, 14. Phytologia Balcanica 16, 415–445.
- VOLOBAEV, P. A., 1991: Genus *Elatine* in Siberia. Sibirskij Biologicheskij Zhurnal 6, 59–64.
- WOHLGEMUTH, T., 1993: Der Verbreitungsatlas der Farn- und Blütenpflanzen der Schweiz (Welten und Sutter 1982) auf EDV. Die Artenzahlen und ihre Abhängigkeit von verschiedenen Faktoren. Botanica Helvetica 103, 55–71.